

Fuzzy Sets and Fuzzy Logic

Fuzzy if-then rules

- General format:


If x is A then y is B

- Examples:


- If pressure is high, then volume is small
- If a restaurant is expensive, then order small dishes
- If a tomato is red, then it is ripe
- If the speed is high, then apply the brake a little

Interpretation of Implication

Common in Fuzzy logic











Implication in traditional logic



A coupled with B

Use the T-norm...



A entails B

- Boolean fuzzy implication (based on $\neg A \vee B$)

$$m_R(x, y) = \max(1 - m_A(x), m_B(y))$$

- Zadeh's max-min implication (based on $\neg A \vee (A \wedge B)$)

$$m_R(x, y) = \max(1 - m_A(x), \min(m_A(x), m_B(y)))$$


- Zadeh's arithmetic implication (based on $\neg A \vee B$)

$$m_R(x, y) = \min(1 - m_A(x) + m_B(y), 1)$$


- Goguen's implication

$$m_R(x, y) = \min(m_B(x) / m_A(y), 1)$$


(a) Zadeh's Arithmetic Rule








(b) Zadeh's Max-Min Rule



(c) Boolean Fuzzy Implication




(d) Goguen's Fuzzy Implication



Building blocks

- Fuzzifier (in the simplest case, turn a measurement into a crisp set)
- Rule base
- Inference engine
- Defuzzifier



Mamdani Systems:


Example 1

- When given are
 - a fuzzy rule $A \rightarrow B$, where A and B are fuzzy sets defined by membership functions $\mu_A(x)$ and $\mu_B(y)$
 - a measurement a for A
- The membership function for $A \rightarrow B$ is defined by
$$\min(\mu_A(x), \mu_B(y))$$
- For a measurement a the membership for y is
$$\min(\mu_A(a), \mu_B(y))$$

Mamdani Systems:


Example 2

- When rules contain multiple conditions, the min is taken over these conditions



Mamdani Systems:

Example 3




Fuzzy observations

- Rule: if x is A then y is B
- Observation: x is A' (fuzzy set)
- Conclusion: y is B' (fuzzy set)

defined as follows:


$$\mu_{B'}(y) = [\bigvee_x (\mu_{A'}(x) \wedge \mu_A(x))] \wedge \mu_B(y)$$
$$= w \wedge \mu_B(y)$$

Graphic Representation




- Rule: if x is A and y is B then z is C
- Fact: x is A' and y is B'
- Conclusion: z is C'

Graphic Representation



Multiple rules, multiple antecedents



Defuzzification rules

- Centroid-of-area

$$z^* = \frac{\int_Z \mu_A(z) z dz}{\int_Z \mu_A(z) dz}$$

- Bisector of area

$$\int_{-\infty}^{z^*} \mu_A(z) dz = \int_{z^*}^{\infty} \mu_A(z) dz$$

- Mean of maximum

$$z^* = \frac{\int_{Z'} z dz}{\int_{Z'} dz}, \quad Z' = \{z \mid \mu_A(z) = \mu^*\}$$

- Smallest of maximum


$$\min_{z \in Z'} z$$

- Largest of maximum

$$\max_{z \in Z'} z$$




range of values
where
membership
is maximal

Mamdani - single input






- X is Small → Y is Small
- X is Medium → Y is Medium
- X is Large → Y is Large

Mamdani - single input



Mamdani - double input



- X is Small and Y is Small
→ Z is negative Large
- X is Small and Y is Large
→ Z is negative Small
- X is Large and Y is Small
→ Z is positive Small
- if X is Large and Y is Large
→ Z is positive Large

Mamdani - double input

