

From lecture 2:

Theorem

Suppose L is a language over the alphabet Σ . If L is accepted by a finite automaton M , and if n is the number of states of M , then

- \forall for every $x \in L$
satisfying $|x| \geq n$
- \exists there are three string u , v , and w ,
such that $x = uvw$ and the following three conditions are true:
 - (1) $|uv| \leq n$,
 - (2) $|v| \geq 1$
- \forall and (3) for all $i \geq 0$, $uv^i w$ belongs to L

[M] Thm. 2.29

Let L be the set of legal C programs.

```
x = main(){{{...}}}
```

[M] E 2.33

Prove the following generalization of the pumping lemma, which can sometimes make it unnecessary to break the proof into cases.

If L can be accepted by an FA,

then there is an integer n

such that for any $x \in L$ with $|x| \geq n$

and for any way of writing x as $x_1x_2x_3$ with $|x_2| = n$,

there are strings u , v and w such that

a. $x_2 = uvw$

b. $|v| \geq 1$

c. For every $i \geq 0$, $x_1uv^iwx_3 \in L$

$$L = \{ a^i b^j c^j \mid i \geq 1 \text{ and } j \geq 0 \} \cup \{ b^j c^k \mid j, k \geq 0 \}$$

- can be pumped, as in the pumping lemma
- is not accepted by FA

[M] E 2.39

Decision problem: problem for which the answer is 'yes' or 'no':

Given ..., is it true that ... ?

*Given an undirected graph $G = (V, E)$,
does G contain a Hamiltonian path?*

*Given a list of integers x_1, x_2, \dots, x_n ,
is the list sorted?*

decidable $\iff \exists$ algorithm that decides

$M = (Q, \Sigma, \delta, q_0, A)$

membership problem $x \in L(M)?$

Specific to M : Given $x \in \Sigma^*$, is $x \in L(M)?$

Arbitrary M : Given FA M with alphabet Σ , and $x \in \Sigma^*$, is $x \in L(M)?$

Decidable, easy

[M] E 2.34

Theorem

The following two problems are decidable

- 1. Given an FA M , is $L(M)$ nonempty?*
- 2. Given an FA M , is $L(M)$ infinite?*

[M] E 2.34

Lemma

Let M be an FA with n states and let $L = L(M)$.

L is nonempty,

*if and only if L contains an element x with $|x| < n$
(at least one such element).*

Theorem

The following two problems are decidable

- 1. Given an FA M , is $L(M)$ nonempty?*
- 2. Given an FA M , is $L(M)$ infinite?*

[M] E 2.34

Lemma

Let M be an FA with n states and let $L = L(M)$.

L is infinite,

*if and only if L contains an element x with $|x| \geq n$
(at least one such element).*

Lemma

Let M be an FA with n states and let $L = L(M)$.

L is infinite,

*if and only if L contains an element x with $|x| \geq n$
(at least one such element).*

Lemma

Let M be an FA with n states and let $L = L(M)$.

L contains an element x with $|x| \geq n$ (at least one such element)


*if and only if L contains an element x with $n \leq |x| < 2n$
(at least one such element).*

- Give 2-state FA for each of the languages over $\{a, b\}$
 - strings with even number of a 's
 - strings with at least one b
- Use the product construction to obtain a 4-state FA for the language of strings with even number of a 's or at least one b
- Investigate which states can be merged

From lecture 1:


Example

$$L_1 = \{ x \in \{a, b\}^* \mid x \text{ ends with } aa \}$$



[M] E. 2.1

Same state, same future



Definition

Let L be language over Σ , and let $x, y \in \Sigma^*$.

Then x, y are *distinguishable* wrt L (*L-distinguishable*),

if there exists $z \in \Sigma^*$ with

$$xz \in L \text{ and } yz \notin L \quad \text{or} \quad xz \notin L \text{ and } yz \in L$$

Such z *distinguishes* x and y wrt L .

Equivalent definition:

$$\text{let } L/x = \{ z \in \Sigma^* \mid xz \in L \}$$

x and y are *L-distinguishable* if $L/x \neq L/y$.

Otherwise, they are *L-indistinguishable*.

The strings in a set $S \subseteq \Sigma^*$ are *pairwise L-distinguishable*, if for every pair x, y of distinct strings in S , x and y are *L-distinguishable*.


Definition independent of FAs

[M] D 2.20

From lecture 1:

Example

$$L_1 = \{ x \in \{a, b\}^* \mid x \text{ ends with } aa \}$$



$$S = \{\Lambda, a, aa\}$$

Theorem

Suppose $M = (Q, \Sigma, q_0, A, \delta)$ is an FA accepting $L \subseteq \Sigma^$.*

If $x, y \in \Sigma^$ are L -distinguishable, then $\delta^*(q_0, x) \neq \delta^*(q_0, y)$.*

For every $n \geq 2$, if there is a set of n pairwise L -distinguishable strings in Σ^ , then Q must contain at least n states.*


Proof...

[M] Thm 2.21

From lecture 1:

Example

$$L_1 = \{ x \in \{a, b\}^* \mid x \text{ ends with } aa \}$$




$$S = \{\Lambda, a, aa\}$$

$$L = \{aa, aab\}^* \{b\}$$

[M] E 2.22

$$L = \{aa, aab\}^* \{b\}$$




[M] E 2.22

Strings with a in the n th symbol from the end

[M] E. 2.24

Strings with a in the n th symbol from the end



[M] E. 2.24

Theorem

For every language $L \subseteq \Sigma^$,
if there is an infinite set S of pairwise L -distinguishable strings,
then L cannot be accepted by a finite automaton.*

Proof...

[M] Thm 2.26

$$Pal = \{x \in \{a, b\}^* \mid x = x^r\}$$